

Lympholyte® Cell Separation Media




Lympholyte®

CEDARLANE's Lympholyte® Cell Separation density gradient centrifugation media has been specifically designed for the isolation of viable lymphocytes from mouse, rat, rabbit, human and other mammalian cell populations. The resulting cell population demonstrates a high, nonselective recovery of viable lymphocytes that are suitable for use as target cells in cytotoxicity and FACS assays and in *in vivo* / *in vitro* functional studies. All products are supplied as sterile liquid with varying densities.

Applications for Lympholyte® include:

- Isolation of viable lymphocytes from lymphoid organs by removal of red and dead cells.
- Removal of dead cells and debris:
 - From lymphocyte suspensions after treatment with antibody plus complement or following cell culture.
 - Following sequential cytotoxicity studies eg. B-cell depletion.
 - From various clonal and hybridoma cell lines.
- Removal of erythrocytes, dead cells and debris From various clonal and hybridoma cell lines. (Lympholyte®-M and Lympholyte®-R)
- Isolation of murine nuclear epidermal cells. (Lympholyte®-M)
- Isolation of mononuclear cells from bone marrow preparations. (Lympholyte®-H)

Lympholyte® Cell Separation Procedure



A. Gently layer the cell suspension or whole blood over top of the Lympholyte® making sure not to mix the two layers.

B. Alternatively, slowly add Lympholyte® to a Pasteur pipette allowing gravity to layer it under the whole blood or cell suspension.

C. Centrifuge the tube at room temperature for indicated time at the indicated speed.

D. Carefully remove the cells at the interface using a Pasteur pipette and transfer into a new centrifuge tube for washing.

Lympholyte® Cell Separation Media

Product	Density	Size	Main Application	Cat. #
Lympholyte®-H Cell Separation Media CE	1.0770 ± 0.001 @ 25°C, g/ml	5x30 mL	Isolation of viable HUMAN lymphocytes from peripheral blood	CL5010
	1.0770 ± 0.001 @ 25°C, g/ml	100 mL	Isolation of viable HUMAN lymphocytes from peripheral blood	CL5015
	1.0770 ± 0.001 @ 25°C, g/ml	6x100 mL	Isolation of viable HUMAN lymphocytes from peripheral blood	CL5016
	1.0770 ± 0.001 @ 25°C, g/ml	500 mL	Isolation of viable HUMAN lymphocytes from peripheral blood	CL5020
	1.0770 ± 0.001 @ 25°C, g/ml	6x500 mL	Isolation of viable HUMAN lymphocytes from peripheral blood	CL5026
Lympholyte®-M Cell Separation Media	1.0875 ± 0.001 @ 25°C, g/ml	5x30 mL	Isolation of viable MOUSE lymphocytes from lymphoid tissue	CL5030
	1.0875 ± 0.001 @ 25°C, g/ml	100 mL	Isolation of viable MOUSE lymphocytes from lymphoid tissue	CL5031
	1.0875 ± 0.001 @ 25°C, g/ml	500 mL	Isolation of viable MOUSE lymphocytes from lymphoid tissue	CL5035
Lympholyte®-Mammal Cell Separation Media For species tested please contact tech@cedarlanelabs.com	1.0860 ± 0.001 @ 25°C, g/ml	5x30 mL	Isolation of viable lymphocytes from peripheral blood of most mammalian species	CL5110
	1.0860 ± 0.001 @ 25°C, g/ml	100 mL	Isolation of viable lymphocytes from peripheral blood of most mammalian species	CL5115
	1.0860 ± 0.001 @ 25°C, g/ml	500 mL	Isolation of viable lymphocytes from peripheral blood of most mammalian species	CL5120
Lympholyte®-Rabbit Cell Separation Media	1.0965 ± 0.001 @ 25°C, g/ml	5x30 mL	Isolation of viable RABBIT lymphocytes from lymphoid tissue	CL5050
Lympholyte®-R Cell Separation Media	1.0940 ± 0.001 @ 25°C, g/ml	5x30 mL	Isolation of viable RAT lymphocytes from lymphoid tissue	CL5040
	1.0940 ± 0.001 @ 25°C, g/ml	100 mL	Isolation of viable RAT lymphocytes from lymphoid tissue	CL5041
	1.0940 ± 0.001 @ 25°C, g/ml	500 mL	Isolation of viable RAT lymphocytes from lymphoid tissue	CL5045
Lympholyte®-poly Cell Separation Media	1.113 ± 0.001 @ 25°C, g/ml	100 mL	Isolation of viable HUMAN polymorphonuclear cells	CL5070
	1.113 ± 0.001 @ 25°C, g/ml	250 mL	Isolation of viable HUMAN polymorphonuclear cells	CL5071
Lympholyte®-1.1	1.100 ± 0.001 @ 25°C, g/ml	500 mL	A high density Lympholyte® solution which can be diluted with PBS (withough Ca ²⁺ /Mg ²⁺). Isolation of pancreatic islet cells prior to expanding them in vitro	CL5095

Lympholyte[®] Cell Separation Media References

Lympholyte[®]-M

Garaycochea JI, Crossan GP, Langevin F, *et al.* (2012) **Genotoxic consequences of endogenous aldehydes on mouse haematopoietic stem cell function.** *Nature.* 489(7417):571-5.

Pulido J, Kottke T, Thompson J, *et al.* (2012) **Using virally expressed melanoma cDNA libraries to identify tumor-associated antigens that cure melanoma.** *Nat Biotechnol.* 18;30(4):337-43.

Y, Marumoto T, Friedmann-Morvinski D, *et al.* (2011) **Transdifferentiation of glioblastoma cells into vascular endothelial cells.** *PNAS.* 108(11):4274-80.

Lympholyte[®]-Mammal

Hoggatt J, Mohammad KS, Singh P, Hoggatt AF, *et al.* (2013) **Differential stem- and progenitor-cell trafficking by prostaglandin E2.** *Nature.* 495(7441):365-9.

Lin WH, Kouyos RD, Adams RJ, Grenfell BT, Griffin DE. (2012) **Prolonged persistence of measles virus RNA is characteristic of primary infection dynamics.** *PNAS.* 109(37):14989-94.

Lympholyte[®]-Rabbit

Cameron CM, Barrett JW, Liu L, *et al.* (2005) **Myxoma virus M141R expresses a viral CD200 (vOX-2) that is responsible for down-regulation of macrophage and T-cell activation *in vivo*.** *J Virol.* 79(10):6052-67.

Lympholyte[®]-R

Bhalli JA, Shaddock JG, Pearce MG, *et al.* (2011) **Report on stage III Pig-a mutation assays using benzo[a]pyrene.** *Environ Mol Mutagen.* 52(9):731-7.

Zhang ZX, Ma Y, Wang H, *et al.* (2006) **Double-negative T cells, activated by xenoantigen, lyse autologous B and T cells using a perforin/granzyme-dependent, Fas-Fas ligand-independent pathway.** *J Immunol.* 177(10):6920-9.

Lympholyte[®]-H

Spinelli E, Caporale R, Buchi F, Masala E, *et al.* (2012) **Distinct Signal Transduction Abnormalities and Erythropoietin Response in Bone Marrow Hematopoietic Cell Subpopulations of Myelodysplastic Syndrome Patients.** *Clin Cancer Res.* 18(11):3079-89.

Colasanti T, Alessandri C, Capozzi A, *et al.* (2012) **Autoantibodies specific to a peptide of β 2-glycoprotein I cross-react with TLR4 inducing a pro-inflammatory phenotype in endothelial cells and monocytes.** *Blood.* 120(16):3360-70.

Pievani A, Belussi C, Klein C, Rambaldi A, Golay J, Introna M. (2011) **Enhanced killing of human B-cell lymphoma targets by combined use of cytokine-induced killer cell (CIK) cultures and anti-CD20 antibodies.** *Blood.* 117(2):510-8.

Lympholyte[®]-Poly

Swystun LL, Mukherjee S, Liaw PC. (2011) **Breast cancer chemotherapy induces the release of cell-free DNA, a novel procoagulant stimulus.** *J Thromb Haemost.* 9(11):2313-21.

Marquez-Curtis L, Jalili A, Deiteren K, *et al.* (2008) **Carboxypeptidase M expressed by human bone marrow cells cleaves the C-terminal lysine of stromal cell-derived factor-1alpha: another player in hematopoietic stem/progenitor cell mobilization?** *Stem Cells.* 26(5):1211-20.

Lympholyte[®]-1.1

Arafat HA, Katakam AK, Chipitsyna G, *et al.* (2007) **Osteopontin protects the islets and beta-cells from interleukin-1 beta-mediated cytotoxicity through negative feedback regulation of nitric oxide.** *Endocrinology.* 148(2):575-84.